



High-fat diet-induced biogenesis of pulmonary exosomes in an experimental rat model

Maryam Shoaran¹ · Mohammad Javad Behmand² · Reza Rahbarghazi³ · Reza Mosaddeghi-Heris⁴ · Mahdi Ahmadi³ · Jafar Rezaie⁵

Received: 8 April 2023 / Accepted: 20 July 2023

© The Author(s), under exclusive licence to Springer Nature B.V. 2023

Abstract

Background High-fat diets (HFD) have recently become a public health concern. We hypothesize that HFD induces exosomes biogenesis in the lung tissue of rat model.

Methods and results Sixteen adult male Wistar rats were fed with HFD or a regular chow diet for 3 months. The histopathological changes in lung tissues were measured by hematoxylin and eosin (H&E) staining. Bronchoalveolar lavage (BAL) was performed to assay exosomes by acetylcholinesterase enzyme (AhCE) activity. Real-time PCR (qPCR) was used to evaluate Rab27-b, Alix, and IL-1 β expression, while the immunohistochemical examination was performed for CD81 expression in lung tissues. In addition, expression of IL-1 β was detected by ELISA. We found pathological alterations in the lung tissue of HFD animals. AhCE activity along with the expression level of Rab27-b, Alix, and IL-1 β was increased in HFD animals ($p < 0.05$). Immunohistochemical staining showed that expression of CD81 was increased in lung tissues of HFD animals compared with the control group ($p < 0.05$).

Conclusion Hence, HFD induced exosomes biogenesis and histopathological changes with IL-1 β expression in rats' lung tissues.

Keywords Exosomes · AhCE activity · HFD · Obesity · Inflammation

Introduction

HFD have become a public health concern due to their role in prompting individuals to obesity and also diabetes [1, 2] as well as inducing overproduction of phosphatidylcholine and insulin resistance [3]. Unusually high levels of cellular phosphatidylcholine lipid impact energy metabolism, which is associated with insulin resistance and obesity [4, 5]. During obesity, oxidative stress and chronic systemic inflammation may affect the function of several organs [6]. In addition, obesity has been considered as important risk factor for several diseases including, asthma, chronic obstructive pulmonary disease (COPD), hypertension, liver and heart diseases, reproductive disorders, and several forms of cancers [7, 8]. An abundant vascular bed in the lungs may make it susceptible to inflammation in an obesity condition, resulting in lung dysfunction. In addition, the accumulation of excess fat tissue in the chest and abdomen regions causes disturbances in the movement of respiratory muscles, especially the diaphragm and intercostal muscles, which causes changes in the functional parameters of the lungs following

Mahdi Ahmadi and Jafar Rezaie contributed equally to this work.

✉ Mahdi Ahmadi
mahdi59866@gmail.com

✉ Jafar Rezaie
Rezaie.j@umsu.ac.ir

¹ Pediatric Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

² Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

³ Stem Cell Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁴ Neurosciences Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁵ Solid Tumor Research Center, Cellular and Molecular Medicine Institute, Urmia University of Medical Sciences, Urmia, Iran

breathing tests [8, 9]. Despite much research, the detailed mechanism behind lung tissue damage in obesity has not been fully determined. Many obesity-related diseases are associated with the hypertrophic and hyperplastic growth of adipocytes which results in adipocyte dysfunction [7, 8]. It is well known that the oxidative stress and pro-inflammatory conditions of the adipocytes may interrupt the secretions of cytokines, adipokines, and other metabolic regulators [10]. Furthermore, under obese conditions, exosomes play a vital role in facilitating the pathogenesis of many diseases [11]. The key roles of adipocyte-released exosomes linking obesity and its comorbidities have been reported in several studies [12, 13]. Exosomes are 30–150 nm membrane-bound vesicles producing from many cells like lung cells, and are present in most body fluids such as bronchoalveolar fluid (BAL) [14]. These vesicles contain many types of proteins, lipids, and RNAs that circulate in almost body fluids, participating in cell-to-cell communication by transferring their contents to nearby or distant cells [15]. The inflammatory condition may cause an alteration in exosomes secretion and cargo [16]. Moreover, there is an association between exosomes secretion and an increase in the level of inflammatory and pro-inflammatory interleukins in patients with COPD and asthma [17]. Therefore, it seems that lung inflammation in obesity may impact exosomes signaling in the lung tissue. Regarding the role of exosomes in obese conditions, in this study, we were looking at whether HFD would impact the secretion and biogenesis of exosomes in lung tissue of the rat HFD model. Understanding exosomes kinetics in lung tissue under HFD conditions may be useful for uncovering the physiological and pathological roles of these exosomes and may offer hints for the progress of exosome-based therapy.

Materials methods

Animals

In this study, we included 16 male Wistar rats weighing between 190 and 10 g, purchased from the animal laboratory unit of Urmia University of Medical Sciences, Urmia, Iran and kept under a 12-h light–dark cycle, temperature control (22 ± 1 °C), with free access to food and water.

Experimental groups

After 1 week of adaptation, rats were randomly allocated into 2 experimental groups, with 8 rats for each group: Control: fed a standard diet; HFD: fed a high fatty pellet diet for three months (Fig. 1).

Diet composition

The essential and standard diet in animal laboratories comprised 11% fat, 22% protein, and 67% carbohydrate. The HFD used in this research contained 32% fat, 20% protein, and 48% carbohydrates for 3 months [18].

Obtaining lung tissues

At the end of the third month, animals were euthanized by anesthetization with xylazine (10 mg/kg) and ketamine (100 mg/kg). The lungs were removed by surgery and immediately, after washing with physiological serum, they were placed in nitrogen and then saved at -80 °C for downstream experimental panels.

Histopathological study


H&E staining was used to investigate the pathological changes caused by HFD in lung tissue. For this purpose, the lung tissues of animals were placed in a 10% formalin solution for fixation. Then, the fixed tissues were placed in different grades of ethanol from 70 to 100% each for 45 min to one hour for dehydration. Next, to prepare tissue blocks, tissue samples were placed in paraffin. Thin sections with a thickness of 4 to 5 μ m were prepared and stained with H&E dye. In this method, hematoxylin turns the nucleus of the cells into purple blue and eosin turns the cytoplasm into pink. Finally, using an optical microscope, pathologic photographs were prepared.

Preparation of BAL

The lungs along with the lower part of the trachea (trachea) were removed surgically. For collecting the BAL, about 1 to 1.5 ml of physiological serum was injected into the lung through a blue branol inserted in the trachea, and then the fluid was instantly collected. This procedure was repeated for each animal a total of 6 times to collect about 5 ml of BAL from each animal.

AhCE activity

AhCE, associated with extracellular vesicles like exosomes, is an important qualitative tool for measuring the exosomes. For this purpose, according to a commercial kit recommendation (BXC080; Iran), 100 μ l BAL samples were incubated with reagent A for 10 min at 37 °C. Then, reagent B was added and after 5 min incubation the absorbance, at five intervals, was calculated at 412 nm

Fig. 1 Timetable and design of the current study

using the recommended formula of a commercial kit as the AhCE activity.

RNA extraction and cDNA synthesis

Total RNA was extracted from the equally homogenized lung tissue using a commercial total RNA extraction mini kit (YektaTajhiz; Iran). UV spectrophotometry technique was used to measure the RNA concentration by NanoDrop 1000 Spectrophotometer (BioTek). The 260 nm/280 nm ratio was also considered a qualitative factor. For reverse transcription, according to the cDNA synthesis kit's recommendation, 1 µg

total RNA was denatured at 70 °C for 5 min, chilled on ice for 5 min and then Transcription Oligo (dT) was added. The samples were incubated with cDNA Synthesis components Mixed at 42 °C for 1 h. The reaction was stopped at 70 °C for 5 min. cDNA samples were stored at – 80 °C for additional amplification.

Real-time PCR (qPCR)

The expression level of Rab27-b, Alix, and IL-1β was determined by qPCR analysis. GAPDH was used as the house-keeping gene. The primers were designed using Oligo-7

Table 1 List of primers

Gene	Primer sequence (5'-3')	
	Forward	Reverse
Rab27-b	GCTGGACCAAAGGGAAGTCA	TCGCCGCACTTGTTTCAAAG
Alix	AGTCGAGAGCCTACTGTGGA	ACTTCTTGAGCTTGGGCCAG
IL-1β	TGGACCTTCCAGGATGAGGACA	GTTTCATCTCGGAGCCTGTAGTG
GAPDH	TTGCCATCAACGACCCCTTCA	AGCACCAGCATCACCCCATTT

software and validated by NCBI database (Table 1). *q*PCR with the cDNA samples was then performed with SYBR green Kit as recommended by the manufacturer (YektaTajhiz; Iran). The PCR reactions were accomplished in a final reaction volume of 14 μ l containing the final concentration of 0.5 μ M of forward and reverse primers in MIC48 instrument (Swiss). Relative expression of genes was measured from Cq values by $\Delta\Delta$ Cq method.

Immunohistochemistry

An immunohistochemical examination was used to evaluate the expression of CD81 in lung tissue samples. In brief, after fixation, tissue samples were embedded in paraffin blocks and mounted on glass slides. Samples were deparaffinized in xylene and then rehydrated in graded alcohol to prepare 5 μ m thick sections. Then, to inhibit endogenous peroxidase activity, sections were treated with hydrogen peroxide (3%) for 10 min and washed with PBS. The slides were then treated with CD81 primary antibody overnight. Next, the slides were incubated with a secondary antibody and then with diaminobenzidine chromogen to visualize the marker. After adding Mayer's hematoxylin and washing with PBS, slides were evaluated under a light microscope and photographed using 5 fields from 3 sections from each rat.

ELISA

To investigate the expression of IL-1 β , we used ELISA according to the manufacturer's recommendations (Rat IL-1 β ELISA kit (ab100768, Abcam)) recommendations. Briefly, Samples were added into the wells for 60 min at room temperature (RT). Then, the wells were washed with PBST solution, and a biotinylated antibody was added. After washing with PBST solution, HRP-conjugated streptavidin was added to the wells. The wells are again washed, and a chromogenic 3,3',5,5'-Tetramethylbenzidine (TMB) substrate solution was added to the wells and kept for 15min for color change in proportion to the amount of IL-1 beta bound.

The reaction was stopped by adding Stop Solution and the color was changed from blue to yellow, and the intensity of the color was measured at 450 nm.

Statistical analysis

The statistical analyses were performed using the program GraphPad Prism 8.0 (California, USA) by T-test. Data were considered statistically significant when *p* value was <0.05. Results obtained from three separate experiments are expressed as the mean $\pm$ SEM.

Results

Histopathological results

As shown in Fig. 2, we observed interstitial pneumonia (black arrow sign) in the lung tissue of the HFD group. No pathological lesion was seen in the control group.


Expression of the IL-1 β gene was up-regulated in HFD rats

The results of both gene expression analysis and ELISA showed that the level of IL-1 β gene expression in the lung tissue of the high-fat diet (HFD) group was significantly increased compared to the control group (*p* < 0.001 and *p* < 0.01; respectively) (Fig. 3A, B), which indicates the inflammatory induction in the lung tissue.

AChE activity increased in exosomes of HFD rats

The AChE activity test showed that the activity of this enzyme in exosomes derived from HFD rats was significantly higher than the control group (*p* < 0.01) (Fig. 4).

Fig. 2 Lung histology in different groups. Healthy lung tissue in the control group (C) and interstitial pneumonia (black arrow sign) in the HFD group are shown. Scale bar was 50 μ m






Fig. 3 Expression of IL-1 β in rats. IL-1 β gene expression in the control group (C) and the high-fat diet group (HFD) (A). The protein level of IL-1 β by ELISA experiment (B). In each group, the results are presented as mean $\pm$ SEM for 8 animals. Student's t-test, ** $p < 0.01$; *** $p < 0.001$ versus the control group




Fig. 4 The level of acetylcholinesterase (AChE) enzyme activity in the lung lavage fluid of the control group (C) and the high-fat diet group (HFD). In each group, the results are presented as mean $\pm$ SEM for 8 animals. Student's t-test, ** $p < 0.01$ versus the control group

Expression of Alix and Rab27-b was increased in HFD rats

QPCR analysis showed that the mRNA levels of Alix and Rab27-b genes were significantly increased in the lung tissue of the HFD group compared to the control group ($p < 0.001$ and $p < 0.01$; respectively) (Fig. 5A). These genes participate in exosomes biogenesis and secretion.

Expression of CD81 was increased in HFD rats

Immunohistochemical staining showed that the level of CD81 protein was increased in the lung tissue of HFD groups (Fig. 5B). This may indicate an increase in the production of exosomes in the lung tissue of the HFD group compared to the control group.




Fig. 5 QPCR analysis for Alix and Rab27b (A). Data showed that expression of Alix and Rab27b was increased in the high-fat diet group (HFD). Immunohistochemical examination for CD81 in the lung tissue samples of the control group (left) and the HFD group (right) (B). Black arrows show CD81 staining in lung tissue. The scale bar was 50 μ m. In each group, the results are presented as mean $\pm$ SEM for 8 animals. Student's t-test, ** $p < 0.01$ and *** $p < 0.001$ versus the control group

Discussion

Exosomes secreted by lung cells into the BAL may contribute to homeostasis and regulate cellular signaling pathways in the lung tissues. Therefore, investigating the exosomes kinetics in the lung tissue can be useful for the diagnosis, treatment, and follow-up of various lung diseases [19, 20]. Previous studies have shown that lung tissue is vulnerable to chronic inflammatory conditions caused by obesity and overweight, however, the main mechanism of lung tissue damage in HFD/obesity has not yet been fully understood. Therefore, this study was conducted to investigate the biogenesis of exosomes in the lung tissue of HFD male rats. The results showed that using an HFD for three months caused inflammation and pathological changes in the lung tissue of male rats along with an increase in IL-1 β expression and the incidence of interstitial pneumonia. These results are consistent with other studies that have shown that obesity and HFD cause inflammation and pathological changes in the lung tissue [21, 22]. Confirmed that obesity and excess adipose tissue contribute to pro-inflammatory conditions [23]. It seems that investigating other lung inflammatory cytokines may uncover the expression level of relevant genes, understanding the connection between HFD pathology and inflammatory cytokines. In the mice model of HFD, it was shown that the level of oxidative stress and inflammatory factors

mediators increased in the lung tissue and alveolar lavage fluid of the animal [21]. We found that rats receiving HFD have a high density of collagen fibers in the lung tissue, possibly due to pulmonary pneumonia and increased collagen expression in the connective tissue, suggesting the incidence of pulmonary fibrosis in these animals [24]. It seems that chronic systemic inflammation in obesity and abundant blood vessels in the lungs are prompting to lung inflammation, pathological changes, and a decrease in its function, however, its exact mechanism has not yet been fully determined. In keeping, we observed that the AChE activity of BAL samples of HFD rats increased, proposing an increase in exosomes/extracellular vesicles production [25]. Then, we were curious if the exosomes biogenesis pathway could be affected by HFD or not. Our finding demonstrated that exosomal markers including Rab27b, Alix, and CD81 were up-regulated in tissue samples of HFD group. Alix and CD81 contribute to exosomes biogenesis and loading of biological cargo into exosomes [26], while Rab27b participates in the intracellular trafficking of exosomes [27, 28]. Our findings would seem to imply that exosomes biogenesis increased in the lung tissues of HFD rats [27, 29]. Previous studies have shown that the levels of circulating fat-derived extracellular vesicles were increased in humans and obese animals, however, decreased following energy restriction [30, 31]. In general, the evidence from this study suggests that an increase in exosomes biogenesis and production was associated with histopathological changes and an increase in IL-1 β expression within the lung tissue of HFD rats. A growing number of studies have found that such pathological conditions as inflammation can modulate exosomes production and cargo loading [32, 33]. Under inflammation, exosomes from affected cells can facilitate pathological conditions [34]. At the same time, alterations in exosomes biogenesis/cargo may serve as disease biomarkers [35]. These data are preliminary and in our opinion, these results widen our knowledge of exosomes biogenesis in HFD condition, however further studies are essential to determine whether HFD condition can alter the contents of exosomes, which can play roles as disease biomarker. In addition, further studies are needed to investigate other inflammatory cytokines to uncover link between exosomes biogenesis and inflammatory condition. In our study, it was not clear how and from which immune cell the exosomes were secreted. Moreover, as the focus of this study was on exosomes biogenesis, there is a possibility that these exosomes carry copies of IL-1 β , which was up-regulated in tissue samples.

Conclusions

Taken together, our research has highlighted inflammation and histopathological changes in rat lung tissue along with an increase in exosome biogenesis after HFD induction. Studying the dynamic of exosomes in the lungs of HFD animals can help us to better understand the pathophysiology and, as a result, innovation of new tools for the diagnosis and treatment of lung-related damage.

Acknowledgements We would like to thank the Clinical Research Development Unit of Zahra Mardani Azari children educational and treatment center, Tabriz university of medical sciences, Tabriz, Iran for their assistance in this research.

Author contributions M-Sh and JR conceived and designed the experiments and analyzed the data. RR prepared the figures. JR and MA wrote the manuscript. R M-H, MA, and MJ-B performed the experiments. All authors read and approved the final manuscript.

Funding This work was approved and supported by a grant from Tabriz University of Medical Sciences (No: IR.TBZMED.VCR.REC.1399.236).

Data availability The data that support the findings of this study are available on request from the corresponding author.

Declarations

Competing interests The authors have no competing interests to declare.

Ethical approval The animal experimental procedures were conducted according to the principles of guidelines for the ethical use of animals in applied studies and approved by the Ethics Committee on Animal Use of Tabriz University of Medical Sciences (IR.TBZMED.VCR.REC.1399.236) in compliance with the ARRIVE guidelines.

Consent for publication Not applicable.

References

1. Canfora EE, Meex RC, Venema K, Blaak EE (2019) Gut microbial metabolites in obesity, NAFLD and T2DM. *Nat Rev Endocrinol* 15(5):261–273
2. Buckley J (2018) Availability of high-fat foods might drive the obesity epidemic. *Nat Rev Endocrinol* 14(10):574–575
3. Pacana T, Cazanave S, Verdianelli A, Patel V, Min H-K, Mirshahi F, Quinlivan E, Sanyal AJ (2015) Dysregulated hepatic methionine metabolism drives homocysteine elevation in diet-induced nonalcoholic fatty liver disease. *PLoS ONE* 10(8):e0136822
4. Kim Y-C, Seok S, Byun S, Kong B, Zhang Y, Guo G, Xie W, Ma J, Kemper B, Kemper JK (2018) AhR and SHP regulate phosphatidylcholine and S-adenosylmethionine levels in the one-carbon cycle. *Nat Commun* 9(1):540
5. Lee S, Norheim F, Gulseth HL, Langleite TM, Aker A, Gundersen TE, Holen T, Birkeland KI, Drevon CA (2018) Skeletal muscle phosphatidylcholine and phosphatidylethanolamine respond

- to exercise and influence insulin sensitivity in men. *Sci Rep* 8(1):6531
6. Blüher M (2019) Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol* 15(5):288–298
 7. Peters U, Suratt BT, Bates JH, Dixon AE (2018) Beyond BMI: obesity and lung disease. *Chest* 153(3):702–709
 8. Dixon AE, Peters U (2018) The effect of obesity on lung function. *Expert Rev Respir Med* 12(9):755–767
 9. Babaei M, Hasanzadeh S, Pirnejad H, Mohebbi I, Hoseini R, Niazkhani Z (2022) Socioeconomic status and severity of traffic accident injuries: a cross-sectional study. *Iran Occup Health* 19(1):380–392
 10. Manna P, Jain SK (2015) Obesity, oxidative stress, adipose tissue dysfunction, and the associated health risks: causes and therapeutic strategies. *Metab Syndr Relat Disord* 13(10):423–444
 11. Kim Y, Son T, Park J, Jun W, Kim OK (2022) Role of exosomes derived from adipose tissue under obese conditions in skeletal muscle and liver cells: commonalities and differences. *Mol Nutr Food Res* 66(23):2200358
 12. Kwan HY, Chen M, Xu K, Chen B (2021) The impact of obesity on adipocyte-derived extracellular vesicles. *Cell Mol Life Sci*. <https://doi.org/10.1007/s00018-021-03973-w>
 13. Maligianni I, Yapijakis C, Bacopoulou F, Chrousos G (2021) The potential role of exosomes in child and adolescent obesity. *Children* 8(3):196
 14. Rezaie J, Feghhi M, Etemadi T (2022) A review on exosomes application in clinical trials: perspective, questions, and challenges. *Cell Commun Signal* 20(1):1–13
 15. Feghhi M, Rezaie J, Akbari A, Jabbari N, Jafari H, Seidi F, Szafert S (2021) Effect of multi-functional polyhydroxylated polyhedral oligomeric silsesquioxane (POSS) nanoparticles on the angiogenesis and exosome biogenesis in human umbilical vein endothelial cells (HUVECs). *Mater Des* 197:109227
 16. Tian Y, Cheng C, Wei Y, Yang F, Li G (2022) The role of exosomes in inflammatory diseases and tumor-related inflammation. *Cells* 11(6):1005
 17. Hough K, Chanda D, Duncan S, Thannickal V, Deshane J (2017) Exosomes in immunoregulation of chronic lung diseases. *Allergy* 72(4):534–544
 18. Hariri N, Thibault L (2010) High-fat diet-induced obesity in animal models. *Nutr Res Rev* 23(2):270–299
 19. Alipoor SD, Mortaz E, Garssen J, Movassaghi M, Mirsaeidi M, Adcock IM (2016) Exosomes and exosomal miRNA in respiratory diseases. *Mediators Inflamm* <https://doi.org/10.1155/2016/5628404>
 20. Almohammedi A, Rahbarghazi R, Keyhanmanesh R, Rezaie J, Ahmadi M (2021) Asthmatic condition induced the activity of exosome secretory pathway in rat pulmonary tissues. *J Inflamm*. 18:1–9
 21. de Souza ABF, Chirico MTT, Cartelle CT, de Paula Costa G, Talvani A, Cangussú SD, de Menezes RCA, Bezerra FS (2018) High-fat diet increases HMGB1 expression and promotes lung inflammation in mice subjected to mechanical ventilation. *Oxid Med Cell Longev* <https://doi.org/10.1155/2018/7457054>
 22. Mafort TT, Rufino R, Costa CH, Lopes AJ (2016) Obesity: systemic and pulmonary complications, biochemical abnormalities and impairment of lung function. *Multidisciplinary Respir Med* 11(1) <https://doi.org/10.1186/s40248-016-0066-z>
 23. Hildebrandt X, Ibrahim M, Peltzer N (2023) Cell death and inflammation during obesity: know my methods, WAT (son). *Cell Death Differ* 30(2):279–292
 24. Wang F, Zuo Z, Chen K, Fang J, Cui H, Shu G, Zhou Y, Chen Z, Huang C, Liu W (2018) Histopathological changes caused by inflammation and oxidative stress in diet-induced-obese mouse following experimental lung injury. *Sci Rep* 8(1):14250
 25. Jabbari N, Feghhi M, Esnaashari O, Soraya H, Rezaie J (2022) Inhibitory effects of gallic acid on the activity of exosomal secretory pathway in breast cancer cell lines: a possible anticancer impact. *BioImpacts* 12(6):549–559
 26. Larios J, Mercier V, Roux A, Gruenberg J (2020) ALIX- and ESCRT-III-dependent sorting of tetraspanins to exosomes. *J Cell Biol* <https://doi.org/10.1083/jcb.201904113>
 27. Ostrowski M, Carmo NB, Krumeich S, Fanget I, Raposo G, Savina A, Moita CF, Schauer K, Hume AN, Freitas RP (2010) Rab27a and Rab27b control different steps of the exosome secretion pathway. *Nat Cell Biol* 12(1):19–30
 28. Hassanpour M, Rezaie J, Darabi M, Hiraifar A, Rahbarghazi R, Nouri M (2020) Autophagy modulation altered differentiation capacity of CD146+ cells toward endothelial cells, pericytes, and cardiomyocytes. *Stem Cell Res Ther* 11(1):1–14
 29. Roucourt B, Meeussen S, Bao J, Zimmermann P, David G (2015) Heparanase activates the syndecan-syntenin-ALIX exosome pathway. *Cell Res* 25(4):412–428
 30. Eguchi A, Lazic M, Armando AM, Phillips SA, Katebian R, Maraka S, Quehenberger O, Sears DD, Feldstein AE (2016) Circulating adipocyte-derived extracellular vesicles are novel markers of metabolic stress. *J Mol Med* 94:1241–1253
 31. Hubal MJ, Nadler EP, Ferrante SC, Barberio MD, Suh JH, Wang J, Dohm GL, Pories WJ, Mietus-Snyder M, Freishtat RJ (2017) Circulating adipocyte-derived exosomal MicroRNAs associated with decreased insulin resistance after gastric bypass. *Obesity* 25(1):102–110
 32. Chaudhuri AD, Dastgheyb RM, Yoo S-W, Trout A, Talbot CC Jr, Hao H, Witwer KW, Haughey NJ (2018) TNF α and IL-1 β modify the miRNA cargo of astrocyte shed extracellular vesicles to regulate neurotrophic signaling in neurons. *Cell Death Dis* 9(3):363
 33. Villarroya-Beltri C, Gutiérrez-Vázquez C, Sánchez-Cabo F, Pérez-Hernández D, Vázquez J, Martín-Cofreces N, Martínez-Herrera DJ, Pascual-Montano A, Mittelbrunn M, Sánchez-Madrid F (2013) Sumoylated hnRNP A2B1 controls the sorting of miRNAs into exosomes through binding to specific motifs. *Nat Commun* 4(1):2980
 34. Tian Y, Cheng C, Wei Y, Yang F, Li G (2022) The role of exosomes in inflammatory diseases and tumor-related inflammation. *Cells*. <https://doi.org/10.3390/cells11061005>
 35. Console L, Scalise M, Indiveri C (2019) Exosomes in inflammation and role as biomarkers. *Clin Chim Acta* 488:165–171. <https://doi.org/10.1016/j.cca.2018.11.009>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.